

# Prevalence and Proportion of Anemia in Pregnant Women Suffering from Malaria

1. Farzana Chang 2. Mir Muhammad Sahito 3. Romela Naz 4. Rabia Shams

5. Riaz Ahmad Sahito

1. Assoc. Prof. of Pathology, LUMHS, Hyderabad 2. Assoc. Prof. of Forensic Medicine, MMC, Mirpur

3. Asstt. Prof. of Forensic Medicine, Sir Syed College of Medical Sciences

4. Prof. of Pathology, PUM&HSW, SBA, 5. Asstt. Prof. Pathology, PUM&HSW, SBA.

## ABSTRACT

**Objective:** To find out the prevalence and proportion of anemia in pregnant women suffering from malaria. The aims & objectives of our study were to evaluate prevalence & proportion of anemia in pregnant women with malaria among the patients visiting the tertiary hospital PUMHS Hospital, Nawabshah, Shaheed Benazirabad so as to give awareness of these complications to doctors to ensure early diagnosis, prevention & prompt treatment of such cases.

**Study Design:** Descriptive observational study.

**Place and Duration of Study:** This study was conducted at Outpatient Department of MMCH Hospital & Department of Pathology from April 2012 to February 2013.

**Materials and Methods:** Study includes total 120 cases of malaria, diagnosed on the basis of clinical & laboratory findings. The hematological complications like anemia, leukocytosis & thrombocytopenia were studied in these patients.

**Results:** Out of 120, 64 patients were anemic. We divided pregnant women into 5 groups, 40 patients were in 16 – 20 years age group and amongst them 55% patients were anemic, 34 patients were in 21 – 25 years age group and amongst them 50% were suffering from anemia, 22 patients were in age group 26-30 & amongst them 22 % anemic, 15 cases were in age group 31 – 35 & out of them 46% were anemic and in age group 36 – 40 years 46 percent were anemic. Overall there were 53% anemic patients in total of 120. According to the occupation status majority were daily wages laborers and farmers and other were housewives. Out of 120 patients most of them were Multigravida. Most of the patients were suffering from anemia, 76% patients had Leucocytosis with Neutrophilia, 74 percent patients had complication of Lymphocytosis, 80 percent of patients had decreased level of monocytes and 60 percent had decreased level of platelets. All the patients were suffering from fever and associated symptoms of chills, sweating, feeling of hotness and coldness, 63 percent had back pain, 45 percent had headache and 23 percent had complication of Splenomegaly. The diagnosis of malaria was made on clinical grounds & confirmed by laboratory findings. The problems of hematological complications as anemia, leukocytosis & thrombocytopenia were detected among the patient of malaria by determining hemoglobin concentration, complete blood picture & urine examination report.

**Conclusions:** P.vivax malaria is most prevalent variety. Anemia, leukocytosis & thrombocytopenia are the most common hematological complications in these patients. It is suggested to ensure

i. Effective malaria control program in country, specially interior Sindh.

ii. Effective measures for prevention, diagnosis & treatment of patients.

iii. Effective health education through electronic & print media.

**Key Words:** Anemia, Pregnant Women, Malaria

## INTRODUCTION

Malaria remains a global health problem with an estimate of 3 billion people at risk of infection in 109 malaria endemic countries. Approximately 250 million cases occur annually leading to approximately one million deaths. 1.2 billion people live in areas with a high risk of malaria (more than 1 reported case per 1000 population per year). There were an estimated 247 million malaria cases among 3.3 billion people at risk in 2006, causing nearly a million deaths, mostly of children under 5 years<sup>1</sup>. In human the sporozoites are transmitted into body by mosquito bite & they first infect the liver cells, then red blood cells & releasing merozoites that mature in male & female

gametocyte in human body. When a healthy female mosquito bites a malaria infected human these gametocytes reach in mosquito's stomach unite together to form zygotes that develop into Oocysts which grow & rupture to release sporozoites & cycle starts again<sup>5</sup>.

According to world malaria report 2011, there were 225 million cases of malaria worldwide & malarial mortality rate reduced by 5% with overall 20-23% throughout world, as an estimated 655000 deaths in 2011 & 781000 deaths in 2010<sup>2</sup>. Although infection due to Plasmodium falciparum is responsible for the greatest overall morbidity and mortality, P.vivax contributes seventy to eighty million new cases to the annual worldwide burden of disease, especially in

temperate regions<sup>3</sup>. Over 40% of world population lives in malaria endemic area including South Asia, India, Pakistan, Bangladesh, Africa, areas of Middle East, Central and South America<sup>4</sup>.

As for hematological complications of malaria are concerned, the anemia, leucocytosis, leucopenia & thrombocytopenia are well recognized complications. The pathogenic mechanisms that cause these complications include haemolysis of parasitized & non-parasitized red blood cells, depression of erythropoiesis, megakaryopoiesis & bone marrow depression, so also immune mediate destruction of thrombocytes caused by releasing of inflammatory cytokines such as tumor necrosis factor & interleukine 1,6 & 10 from the activated macrophages in malaria<sup>6</sup>. The factors contributing to the severity of hematological complication are malaria immunity, endemicity, hyperparacytemia, hemoglobinopathies, G6PD deficiency nutritional status & co-morbidity of HIV & dengue fever<sup>7</sup>.

The clinical presentation of malaria caused by all species of malaria parasite resembles & includes fever, headache, body ache, joint pain, chills, sweating & vomiting. These symptoms appear within 1 – 2 weeks after bite of infective mosquito<sup>8</sup>.

*Pl. falciparum* originally was discovered in gorillas caused sever malaria especially among the children producing anemia, respiratory distress, hepatosplenomegaly & finally renal failure or black water fever & cerebral malaria with retinal changes (whitening) which may be a useful clinical sign that distinguish malaria from other causes of fever<sup>9,10</sup>. Although blood is most frequently used to make a diagnosis of malaria by microscopic examination of Giemsa stained blood films but saliva & urine have been investigated as an alternative non invasive techniques<sup>11</sup>. The rapid malaria diagnosis test & polymerase chain reaction have been discovered recently to be used for diagnosis of malaria. These tests are not implemented in poor country like Mozambique due to their high cost. Often the history of fever is taken as the indication to treat malaria in these countries<sup>13,14,15</sup>.

## MATERIALS AND METHODS

This Observational descriptive study was conducted at outpatient department of MMCH Hospital & Department of Pathology from Apr 2012 to Feb 2013. Total 120 pregnant patients visiting OPD were included in the study and patient having any morbidity were excluded from the study. The prevalence of malaria with reference to age, sex, area of residence & clinical findings were recorded. For the laboratory diagnosis of malaria blood samples in test tube containing EDTA were sent to pathology department. Thick & thin blood smears were made on the clean glass slides from the EDTA mixed blood, &

examined under microscope for detection of various developmental stages of malaria parasites after staining with Giemsa stains. The Hb concentration, CBC including RBC, TLC, DLC & platelet counts were determined by a hematological analyzer (Sysmac). The ESR & ICT were also done from the same blood sample.

## RESULTS

Out of 120 pregnant patients, 40 were from urban and 80 pregnant women were from rural life. Out of 40, 19 patients were anemic and 21 were non-anemic while out of 80 rural patients 50% were anemic (Table No. 1). We divided pregnant women into 5 groups, 40 patients were from 16-20 year (anaemic patient were 22), 34 patients were from 21-25 year (50 percent women were suffering from anemia), 20 patients were in age group 26-30 years (11 patients were anaemic), 15 patients were between 31-35 (7 patients were anaemic) and 11 were in 36-40 years group (7 were anaemic). Overall 64 patients were anemic in total 120 cases (Table No.2). Regarding occupational status, majority were daily-wages laborers and farmers and rest were housewives (Table No.3). Out of 120 patients majority were multigravida (Table No.4). All patients were suffering from anemia but 76% had Leucocytosis with Neutrophilia, 74 percent patient had complication of Lymphocytosis, 80 percent of patient had decreased level of monocytes and 60 percent had decreased level of platelet (Table No. 5). All the patients were suffering from fever and associated symptoms of chills, sweating, feeling of hotness and coldness, 63 percent had body pain, 45 percent had headache and 23 percent had complication of Splenomegaly (Table No. 6).

**Table No. 1: Distribution of case with reference to residence**

| Place of residence | Anemic      | Non – Anemic |
|--------------------|-------------|--------------|
| Urban 40           | 19 (47.5%)  | 21 (52.5%)   |
| Rural 80           | 45 (56.25%) | 35 (43.75%)  |
| Total 120          | 64(53.33)   | 56(46.67)    |

**Table No. 2: Distribution of case with reference to age**

| Age related group | Total case | Anemic     | Non – Anemic |
|-------------------|------------|------------|--------------|
| 16-20 yrs         | 40         | 22 (55%)   | 18 (45%)     |
| 21-25 yrs         | 34         | 17 (50%)   | 17 (50%)     |
| 26-30 yrs         | 20         | 11 (55%)   | 9 (45%)      |
| 31-35 yrs         | 15         | 7 (46.66%) | 8 (53.33%)   |
| 36-40 yrs         | 11         | 7 (46.66%) | 4 (36.66%)   |
| Total             | 120        | 64(53.33)  | 56(46.67)    |

**Table No. 3: Distribution of case with reference to occupation**

| Sr. No. | Occupation group   | Total case | Anemic         | Non – Anemic  |
|---------|--------------------|------------|----------------|---------------|
| 1       | Housewife          | 15         | 10<br>(66.66%) | 5<br>(33.33%) |
| 2       | Farmer             | 30         | 17<br>(56.66%) | 13<br>(43.3%) |
| 3       | Daily-wages worker | 75         | 60 (80%)       | 15 (20%)      |

**Table No. 4: Distribution of case with reference to parity**

| S. No. | Parity          | Total case | Anemic         | Non – Anemic   |
|--------|-----------------|------------|----------------|----------------|
| 1      | Primigravida    | 35         | 18<br>(51.42%) | 17<br>(48.57%) |
| 2      | Multigravida    | 58         | 40<br>(68.96%) | 18<br>(31.04%) |
| 3      | Third trimester | 27         | 18<br>(66.66%) | 9<br>(33.33%)  |

**Table No. 5: Distribution of case with reference to haematological complications**

| Sr. No. | Haematological Complications   | No. of patients | Percentage |
|---------|--------------------------------|-----------------|------------|
| 1       | Anemia                         | 120             | 100.00     |
| 2       | Leucocytosis with Neutrophilia | 92              | 76.60      |
| 3       | Lymphocytosis                  | 89              | 74.16      |
| 4       | Decrease in Monocytes          | 95              | 80.00      |
| 5       | Thrombocytopenia               | 72              | 60.00      |

**Table No. 6: Distribution of case with reference to clinical findings**

| Sr. No | Clinical Findings                                                           | No. of Patient | Percentage |
|--------|-----------------------------------------------------------------------------|----------------|------------|
| 1      | Fever                                                                       | 120            | 100%       |
| 2      | Associated Symptoms like Chills, Sweating, feeling of Coldness and hotness. | 120            | 100%       |
| 3      | Bodyache                                                                    | 76             | 63.33%     |
| 4      | Headache                                                                    | 55             | 45.83%     |
| 5      | Pallor                                                                      | 80             | 66.66%     |
| 6      | Splenomegaly                                                                | 28             | 23.33%     |

**Table No.7: Distribution of cases with reference to haematological parameters(Hb & ESR)**

| Sr. No. | Haematological parameters         | No. of Patient | Percentage |
|---------|-----------------------------------|----------------|------------|
| 1       | HB<br>5.5 – 11.5 g/dl $8.5 \pm 3$ | 88             | 73.30      |
| 2       | ESR<br>40 – 110mm 37.5 – 72.5     | 120            | 100.00     |

**Table No. 8: Distribution of cases with reference to hematological parameters determined by haematological analyser (Sysmac)**

| Sr. No. | Haematological parameters                         | No. of patients | Percentage |
|---------|---------------------------------------------------|-----------------|------------|
| 1       | TLC: 6500-25000/cumm<br>Mean= $8750 \pm 1625$     | 90              | 75%        |
| 2       | RBCs: 2.4-4.5m<br>Mean= $3.5 \pm 1.0$             | 67              | 55.8%      |
| 3       | DLC: Eutrophils<br>67-85%<br>Mean= $80.5 \pm 5.5$ | 92              | 76.6%      |
| 4       | Lymphocytes<br>10-14%<br>Mean= $11 \pm 3$         | 87              | 72.5%      |
| 5       | Monocytes: 10-18%<br>Mean= $14 \pm 4$             | 96              | 80.0%      |
| 6       | Eosinophils: 2-4%<br>Mean= $3 \pm 1$              | 120             | 100%       |

## DISCUSSION

The malaria mortality rate in 2010 was 50, 56 & 86 persons per 100,000 populations in 3 African countries as reported by Guardian news London<sup>16</sup>. On the celebration of malaria day on 25 April 2011, the map of *Plasmodium falciparum* malaria & endemicity in 2010 & several other maps related to malaria were published in Asia, Africa & America as noted by Gething et al<sup>17</sup>.

Malaria was endemic in broadband around the equator, in areas of America, many part of Asia and much of Africa, & 85 – 90% of malaria mortality due to the *P. falciparum* occurred among the children & pregnant women in Sub-Saharan Africa<sup>18</sup>.

Malaria was prevalent throughout the human history & one in every two people had been died due to malaria & its complications. Atlas projects founded by Wellcome UK Trust to rectify malaria mapping providing more contemporary & robust means by which the current & future malaria disease burden was assessed as reported by Hay et al<sup>19</sup>.

The malaria superimposed with HIV infection is increasing a person's susceptibility to malaria infection as stated by Abu Redid et al<sup>20</sup>.

The female anopheles mosquitoes such as *A.culicifacies* & *A.sephenisi* were common in Pakistan that transmitted malaria as recorded by national Malaria control program 2006<sup>21</sup>. While prevalent rate of *P. Vivax* malaria was two times higher than *P. falciparum* in interior Sindh as well as in Baluchistan Pakistan, Mehmood et al<sup>22</sup>, Nizamani et al<sup>23</sup> & Yasin Zai et al<sup>24</sup>.

The hematological complications in malaria such as anemia, Leucocytosis with Neutrophilia & thrombocytopenia were stated by Khalid et al<sup>25</sup> & age as a risk factor for thrombocytopenia & anemia in children treated for acute uncomplicated *falciparum* malaria was found by Adedapo et al<sup>26</sup>. The high

concentration of interleukin-10 associated with thrombocytopenia in falciparum malaria was demonstrated by Casal-Pascual et al<sup>27</sup>. The hemozoin containing leukocytes (HCL) such as monocytes & neutrophils seen in blood films on patients with malaria were negatively correlated with severity of malaria in Ugandan children as stated by Mujuzi et al in his study<sup>28</sup>.

In contrast to this study, Hunched et al studies stated that presence of HCL in blood films of patients with malaria were used as reliable diagnostic but less prognostic tools<sup>29</sup>. Both studies stated that hemozoin is brownish black malaria pigment that digests product of hemoglobin produced by phagocytosis of parasitized & none parasitized RBCs by monocytes & neutrophils. Casal – Pascual et al demonstrated that suppression of erythropoiesis produced severe malarial anemia that is associated with hemozoin containing leukocytes in peripheral blood<sup>30</sup>.

The hematological complications of malaria evaluated by hematological parameters to differentiate malaria from other acute febrile illness were studied by Danish et al<sup>31</sup>. They stated that severe anemia & thrombocytopenia determined by hemoglobin & platelets counts less than 5g/dl & 5000/cumm respectively in these patients had more specific, sensitive & predictive values to diagnose malaria.

## CONCLUSION

*P. vivax* malaria is most prevalent variety. Anemia, leukocytosis & thrombocytopenia are the most common hematological complications in these patients. It is suggested to ensure

- i. Effective malaria control program in country, specially interior Sindh.
- ii. Effective measures for prevention, diagnosis & treatment of patients.
- iii. Effective health education through electronic & print media.

## REFERENCES

1. Rehman A. Malaria: The trend at Tehsil Hospital Liaquatpur. Prof Med J 2005;12(3):322-326
2. World Health Organization; World malaria report 2011.
3. Rehman A. The passive case detection of malaria at Tehsil Liaquatpur. Prof Med J 2006;13(1): 32-24.
4. Hayat AS, Siddique MS, Sheikh N, Muqetullah. Thrombocytopenia: frequency and degree in patients with falciparum malaria. Prof Med J 2011; 18(1):75-79.
5. Palaisa M. Life cycle of Malarial Parasites. Med J Thera Africa 2008; 2(3):227-8.
6. Douglas JP, Gregoge DP, Parkash K, Jane BH, Jhon Michael. Several Malarial anemia; Innate

immunity and pathogenesis. Int J Bio Sci 2011; 7(9):1427-42.

7. Maina R, Erhart LM, Chuanak N. Hematological complication of malaria. Malaria J 2010; 9:1-19.
8. Rasheed A, Saeed A, Khan SA. Clinical presentation in acute malaria by all species of plasmodium. JPMA 2009; (4):220-22.
9. Liu WY, Li GHL, Rudicell RS. Origin of human malaria Parasite Plasmodium Falciparum in gorillas. Nature 2010;467:4205.
10. Maude RJ, Hassan MU, Beare NAV. Severe Retinal whitening in adult in cerebral malaria. AM J Trop Med Hyg 2009; 80(6):881.
11. Krafts K, Hempelmann E, Oleksyn B. Detection of malarial parasite by Geimsa stained blood films. Biotechm 2011;86(1):735 .
12. Sutherland CJ, Hallett R. Detecting Malarial parasites outside the blood. J Infect Dis 2009; 199(11):1561-3.
13. Malkar MT, Piper RC. The rapid malaria diagnostic test. AM J Trop Med Hyg 2009; 81(6):921-6.
14. Mens PF, Schoone GJ, Kager PA, Schalling HD. Detection and identification of human Plasmodium species with real time quantitative nucleic acid sequence based amplification. Malaria J 2006; 5(8):80.
15. Hame JC., Barnish G, Mangal T, Armazio I, Street E, Beats I. Household cost of malaria over diagnosis in rural Mozambique. Malaria J 2008; 7: 33.
16. Provost and Claire Wold Malaria Day; which countries are the hardest hit? Get the full data, the guardian (London) 25 April 2011.
17. Gething PW, Patil AP, Smith DL. A new malaria World Map; Plasmodium Falciparum endemicity in 2010. Malaria J 2011;10(1):378.
18. World Malaria Report WHO 2008.
19. Hay SL, Snow RW. The Malaria Atlas Project; Developing Maps of Malaria Risk. Plo S Med 2006;(12):473.
20. Raddad AL, Patniak P, Kublin J. Dual infection with HIV and malaria fuels the spread of both diseases in Sub-Saharan Africa. Sci 2006; (5805): 1603-6.
21. MPC. District wise epidemiology data of malaria control program, Balouchistan, Pakistan, Islamabad. Malaria Control Program 2006.
22. Mahmood KH. Malaria in Karachi and other areas in Sindh. PAF Med J 2005;55:345-8.
23. Nizamani A, Kakar NA, Khushk I. A burden of malaira in Sindh, Pakistan a two years surveillance report JLUMHS 2006;5:76-82.
24. Zia YMI, Khel KSJK. Incidence of malaria in central areas of Balouchistan; Mastung and Khuzdar. Rawal Med J 2007;32:176-8.

25. Taha K, EL-Dein SZ, Idress M, Malout G, Ghasan Baidass. Hemotological complications of malaria. Kuwait Med J 2007;39(3);262-7.
26. Adedapo AD, Falade CO, Kotila RT, Ademowo GO. Age as a risk factor for Thrombocytopenia and anemia in children treated for acute uncomplicated falciparum malaria. J Vector Borne Dis 2007; 44:266-71.
27. Casal-Pascual C, Kai O, Newton CR, Peshu N, Roberts DJ. Thrombocytopenia in falciparum malaria is associated with high concentration of IL-10. AM J Trop Med Hyg 2006;75:434-6.
28. Mujazi G, Magsmbo B, Okech B, Egwang TG. Hemozoin containing leucocytes (HCL) are negative correlates of protection against severe and complicated malaria in Ugandan children. Am J Trop Med Hyd 2006;74:724-9.
29. Hanscheid T, Egan TJ, Grobusch MP. Hemozoin from melatonin pigment to drug target, diagnostic tool, and immune modulator. Lancet Infect Dis 2007;7: 675-85.
30. Casal-Pascual C, Kai O, Cheung Jo, Hillenes S, et al. Suppression of erythropoiesis in malarial anemia associated with hemozoin concentration leucocytes in blood. Blood 2006; 108:2569-77.
31. Dinesh R, Viral P, Amarjeet AK, Vinod DP, Devangi DP. Haematological Parameters differentiating acute malaria from other acute febrile disease including dengue fever. Ind J Pathol Micro 2009;32(2):185-8.

**Address for Corresponding Author:****Dr. Farzana Chang,**Associate Professor Pathology,  
Liaquat University of Medical & Health Sciences,  
Jamshoro.

Email: drfarzana1@hotmail.com

Electronic Copy