

Study of UTI Causing Organisms in Patients Visiting Tertiary Care Hospitals

Muhammad Amjad, Muhammad Shafiq and Muhammad Safdar

ABSTRACT

Objective: To assess the frequency of UTI causing organisms and to determine the frequency in both the genders.

Study Design: Prospective / cross sectional study

Place and Duration of Study: This study was conducted at the tertiary care hospital from July 2015 to June 2016.

Materials and Methods: The patients presented at the urology department with complaints of urinary tract infection aging 21-70 years were enrolled with simple purposive sampling in the study. The majority of the patients got enrolled were aging 21-35 years. The data was collected with the help of preformed questionnaire.

Results: 340 patients were enrolled in the study 202 females and 138 males by purposive sampling. The urine sample was collected and sent to hospital laboratory for culture and examination. most of cases with positive culture with identified organisms had klebsiellapneumonia, Staphylococcus sp., Escherichia Coli, Pseudomonas, Proteus SP and Candida's.

Conclusion: It is concluded from the results of the study that E. coli is the most common organism causing UTI in both males and females.

Key Words: Urinary tract infections, isolated microorganisms, Patients.

Citation of articles: Amjad M, Shafiq M, Safdar M. Study of UTI Causing Organisms in Patients Visiting Tertiary Care Hospitals. Med Forum 2018;29(9):65-67.

INTRODUCTION

The Urinary tract infection is the condition having multiple clinical symptoms that are clinically positive ranging from asymptomatic bacteriuria that may lead to kidney infections¹. It is a serious health concern and is the 2nd most common disease of the body affecting the millions of the people and is a serious public health problem. According to an estimate, 150 million cases of urinary tract infections are diagnosed per annum². The urinary tract infections are the most common bacterial infections being presented to clinicians on routine basis in the developing countries. A patient can be affected by UTI in any time in his life, in all ages including childhood and old age in both genders³. The clinical symptoms of urinary tract infections include urinary pain or burning sensation while urinating, cola or vivid pink color and blood signs, more frequent as well as urgency to urinate. If the urinary bladder is filled with urine, the patient may sense pain in the pubic bone, infemale's pelvic pain and in male's rectal pain and smell of cloudiness, even after the disease is softened the bladder still reacts sensitively⁴. In infants and children, the clinical symptom of the urinary tract Infection can be fever, vomiting, more sleep, signs of

malnutrition, or jaundice. Newly manifesting urinary incontinence may occur in elderly children⁵. The normal microbial flora is the habitant of parts of the body or reside in it. Normally bacteria are also found in vagina and urethral area that rarely cause disease in healthy people. After the child is born, the peri-urethral area gets normal habitants of bacteria to counter the UTI pathogens. when the patient is given antibiotics for any other disease, the normal flora is destroyed and the risk of the UTI is increased. the lactobacillus species reside in the vagina and there are responsible for the specific odour of the vaginal are, they produce lactic acid and some produce hydrogen peroxide, during mensuration the normal vaginal flora is disturbed and hence the risks of UTI are increased⁶. Almost 95% of the cases of UTI are due to bacteria that normally is habitant of the opening of urethra and then migrate to bladder. Most of the infections are due to retrograde rise in the number of bacteria from the faecal plexus via urethra to the bladder and then to the kidney, and it is more seen in women due to shorter urethra. The prevalence of all the UTIs are increasing in origin and the cause is the flora off the gastrointestinal tract which occupies per urethral area⁷. The increased incidence of UTI is in females as compared to males, the reason may be due to trauma during sexual intercourse and chronic pathogens in the perennal skin and in males the pathogens are found in the capsule. In case of infants, the congenital anomalies are the main reasons for UTIs. the prevalence of females compared to males, the more severe pollution during sexual intercourse, urethral massage, and even chronic pathogens present in perennal skin. In males, the organisms are often come

Department of Urology Kidney Center BVH Bahawalpur.

Correspondence: Dr. Muhammad Amjad, Senior Registrar,
Department of Urology Kidney Center BVH Bahawalpur.
Contact No: 0334-7090802
Email: dramjad284@hotmail.com

Received: January, 2018;

Accepted: April, 2018

under the capsule. And in elderly, the obstruction due to prostate enlargement causes partial emptying of the bladder and increases the risk of the UTI⁸. It is really important to treat the disease as soon as possible to reduce the risks for long term morbidity and mortality.⁹ The proper treatment requires the understanding of etiology and the bacterial factors and role of host. proper diagnosis and management in most of the cases result in the success in resolution of the infection¹⁰.

MATERIALS AND METHODS

It is a cross sectional study done at a Urology Kidney Center BVH Bahawalpur. Duration of the study from July 2015 to June 2016. In the study, 340 cases were enrolled. Patients presenting at the Urology department with symptoms of urinary tract infections were enrolled in the study by the purposive sampling. Patient's urine was collected and sent for culture and examination. The Data was taken with the help of preformed questionnaire. the data analysis was done by spss20. Data management was done properly keeping the filled questionnaires and all other relevant documents in safe lockers and putting the data on daily in SPSS to analyze statistically.

RESULTS

All of the 340 patients having UTI and positive urine culture were enrolled in our study. There are 202 (64%) females having positive urine culture and 108 (36%) males having positive urine culture.

Table No.1: Percentage of microorganisms with sex distribution

Microorganism	Frequency (Percentage)	Sex	
		Female	Male
Klebsiellapneumonea	64.6 (19%)	26 (8.6%)	34 (10.3%)
Escherichia Coli	132 (39%)	78 (26%)	44 (13%)
Staphylococcus sp.	17 (5.7%)	10 (3.6%)	9 (3%)
Proteus sp.	15 (5%)	9 (3%)	6 (2%)
Candida	7 (2.3%)	2 (0.67%)	5 (1.6%)
Pseudomonas	99 (29%)	66 (22%)	24 (7%)

Cases enrolled in our study aged 21 to 70 years. And bacteria identified of urine culture are as: Klebsiellapneumonea, Escherichia Coli, Candida and Pseudomonas. Staphylococcus sp., Proteus sp percentage and frequency of bacteria is given in the table 1. The most highly prevalent of the microorganisms is E. coli and the least frequent is Candida. The most frequent of the pathogen among females was recorded to be of E. coli (26%) and same is highest among males (12%).

DISCUSSION

Urinary tract infections pose serious challenge for the community and hospitals. According to surveys conducted, UTIs are the commonest bacterial disease reported in emergency and OPD patients. The hundred thousand of hospitalizations per annum are due to UTIs, 1.1 million emergency patients and 7 million OPD patients, causing a huge financial burden, as per an estimate 1.6 billion are being paid per year due to UTI.¹¹ There is variability in the results of prevalence of UTI depending on the age and sex of the cases¹². Urinary tract infections are More common among females than in males, 33% of the females are affected by uti in life time for at least once¹³.

In this study, the highly prevalent age for this disease is 21-35 years of age. and more female patients than males. 340 patients were enrolled in our study and 202 were females and 138 were males. The prevalence is near to the frequency reported by Ahmed and Avasarala i.e 12.7%¹⁴ but it is higher than the study done by Singh MM et al. who reported it to be 4.2% UTI¹⁵. In Bangladesh, patients usually visit a doctor after experiencing severe complications for a disease condition. So, in a studies done by Bashar et al. and Rahman et al. higher frequency of UTI is reported i.e. 27% and 24.14%^{16,17}. In our study, UTI was most frequently found in age group 21-35 years, the results are similar to the other studies done. The incidence of uti is higher in females than males, females have marked high percentage of the urine culture positivity i.e 90.15% and males 9.85% of total cultures sent for examination. Out of 192 strains, 175 were females and 17 were males Similarity in observations has been seen in the studies done by Astal et al.²⁶, Khalifa et al¹⁸, Bangladesh (Begum et al., 16.4% of female urinary tract infections in Dhaka).

The culture results usually show the Gram-negative intestinal bacteria as culprit, that mostly cause urinary tract infections, e.g E. coli, Klebsiella spp. And Proteus species¹⁹. The most commonly involved bacteria in UTI (80-85%) are E. coli. 30. In our study, the Escherichia coli had the largest of the groups with a frequency of 39%, then comes Pseudomonas sp 29% and then Klebsiellasp 19% and Proteus, Staphylococcus sp, and Candida were below 10%. Other researchers (Basar et al. and Saber et al. also indicated higher correlation of E. coli (66.67% and 77.8% patients respectively) in patients of UTI^{20,21}. Another study done in 2014 in Lahore shows the frequency of UTI is with the highest prevalence of E-coli (80%) then followed by Staphylococcus aureus (9.4%), Pseudomonas species (5.2%) and Proteus species 5.4%. In most of the studies conducted Escherichia coli has been the most prevalent organism.²²

CONCLUSION

The Urinary tract infections are more frequent in females as compared to males and enteric bacteria *E. coli* is the most prevalent pathogen among both males and females. Identification and diagnosis of microorganisms are important and very helpful for physicians to decide and treat these patients accordingly.

Author's Contribution:

Concept & Design of Study: Muhammad Amjad
 Drafting: Muhammad Shafiq
 Data Analysis: Muhammad Safdar
 Revisiting Critically: Muhammad Amjad, Muhammad Shafiq
 Final Approval of version: Muhammad Amjad

Conflict of Interest: The study has no conflict of interest to declare by any author.

REFERENCES

1. Tanagho, Emil A, Mcaninch, Jack W. Smith's General Urology: United States of America. McGraw Hill companies Inc; Bacterial Infections of the genitourinary tract 2004;2:203-227.
2. Owa JA. Urinary tract infections in children, Paediatrics and child health in tropical region 2007;2(2).
3. Tessema B, Kassu A, Mulu A, Yismaw G. Predominant isolates of urinary tract pathogens and their antimicrobial susceptibility patterns in Gondar University Teaching Hospital North West Ethiopia. *Ethiop. Med J* 2007;(1):2007:61-67.
4. Car J. Urinary tract infections in women: diagnosis and management in primary care. *BMJ* 2006;(332): 94-97.
5. Bhat RG, Katy, TA, Place FC. Pediatric urinary tract infections. *Emergency medicine clinics of North America* 2011;29(3):637-653.
6. Vasquez A, Jakobsson T, Ahme S, Forsum U, Molin G. Vaginal Lactobacillus Flora of Healthy Swedish Women. *J Clin Micro* 2002;40(8): 2746-2749.
7. John ED and Michel L. Urinary tract infections during pregnancy. *J Am Fam Physician* 2006; 61(3):713-720.
8. Starr C, Taggart RC. Biology the unit and diversity of life. Wards worth publishing Co. Belmont co 2002;(3):509-533.
9. Quigley. Diagnosis of Urinary Tract Infections in children. *Curr Opin Pediatr* 2009;21(2):124-128.
10. Masson P, Matheson S, Webster AC, Craig JC. Meta-analyses in prevention and treatment of urinary tract infections. *Infect Dis Clin North Am* 2009;(23):355-385.
11. Joseph TD. Urinary Tract Infections and Prostatitis. In: Wells BG, Schwinhart TL, Hamilton CW, editors. *Pharmacotherapy Handbook*. 7th ed. New York: McGraw-Hill;2008. p.493-503.
12. Kosokai NY, Kumaoto T, Hirose N, Tanka Y, Ltikichi S, Sigeta Y, et al. Comparative studies on activities of antimicrobial gaunt against causative organisms isolated from urinary tract infection of 1987.11 Background of patients. *Japan J Antiriot* 1990;43:954-67.
13. Palac DM. Urinary tract infection in women. A physician's perspective 1986;17-25.
14. Ahmed SM, Avasarala AK. Urinary tract infections (UTI) among adolescent girls in rural Karimnagar district, AP - K.A.P. STUDY. *Ind J Prev Soc Med* 2008;39(12):67-70.
15. Singh MM, Devi R, Garg S, Mehra M. Effectiveness of syndromic approach in management of reproductive tract infections in women. *Indian J Med Sci* 2001;55(4):209-214.
16. Bashar MA, Ahmed MF, Rahman SR, Gomes DJ. Distribution and Resistance Trends of *Escherichia coli* from Urinary Tract Infections Isolated in Dhaka City. *Ban J Med Sci* 2009;15(2):93-98.
17. Rahman F, Chowdhury S, Rahman MM, Ahmed D, Hossain A. Antimicrobial Resistance Pattern of Gram-negative Bacteria Causing Urinary Tract Infection. *S J Pharm Sci* 2009;2(1):44-50.
18. Bhowmick B K, Rashid H. Prevalence and antibiotic susceptibility of *E coli* Isolated from Urinary tract infections (UTI) in Bangladesh. *Pak J Biol Sci* 2004;7 (5):717-720.
19. Astal Z, Sharif F, Manama A. Antibiotic resistance of bacteria associated with community-acquired urinary tract infections in the southern area of the Gaza Strip. *J Chemother* 2002;14:259-64.
20. Khalifa BHA, Khedher M. Epidemiological study of *Klebsiella* spp. uropathogenic strains producing extended-spectrum betalactamase in a Tunisian University Hospital, 2009. *Pathol Biol (Paris)* 2012;60: e1-5
21. Begum N, Mamoon ABA, Hossain M, Begum N, Chowdhury SA, Rahman MF. UTI among female workers in a selected garment industry of Dhaka city: A cross sectional study. *The ORION Med J* 2006; 23:325-327.
22. Bova JG, Potter JL, Arevalos E, Hopens T, Goldstein HM, Radwin HM. Renal and perirenal infection: to the role of computerized tomography. *J Urol* 1985; 133:375-8.
23. Nicolle LE. Resistant pathogens in urinary tract infections. *J Am Geriatr Soc* 2002;50: S230-5